

PEPI FINDER

This is an index to the statistical procedures provided by WINPEPI (PEPI-for-Windows) programs. It may be used to find what program and module should be used, and as an index to the manuals. “**COMPARE2 A, B**”, for example, refers to modules **A** and **B** of **COMPARE2**.

The WINPEPI programs are: **COMPARE2** (comparison of two groups or samples) v.1.57
DESCRIBE (descriptive epidemiology) v.1.69
ETCETERA (miscellaneous) v.1.70
LOGISTIC (multiple logistic regression) v. 1.15
PAIRSetc (comparison of matched observations) v.1.43
POISSON (Poisson regression analysis) v.1.11
WHATIS (“ready reckoner” utility program) v.4.21.

The list includes procedures that are provided only by PEPI DOS programs (which are shown in *ITALICS*).

Updated 30th June, 2007.

2 x 2 table, **COMPARE2 A, B**
2 x 2 table, case-control study, **COMPARE2 C**
2 x 2 table, for paired data, **PAIRSetc A**
2 x k table, **COMPARE2 F1**
2 x k table, ordered categories, **COMPARE2 F2, G**
95% limits of agreement, **PAIRSetc D1, D2, D3, L2**

A
AC₁ statistic, **PAIRSetc A**
Adjusted rank correlation test for skewed funnel plot, **COMPARE2 A, B, C, D, I**
Alpha (Cronbach’s reliability-consistency coefficient), **ETCETERA D**
Analysis of covariance, for comparing changes in two groups, **PAIRSetc D1, D2**
Annual rate of increase or decrease, **DESCRIBE B**
ANOVA (analysis of variance) table, **COMPARE2 E, PAIRSetc D1, H, I1, I2, L2**
Attributable fraction, **COMPARE2 B, C, G, PAIRSetc A**

B
BAK (bias-adjusted kappa), **PAIRSetc A, B, C**
Bartholomew’s test for trend, **DESCRIBE B**
Bartlett’s test for comparing two variances, **COMPARE2 H2**
Basic reproductive ratio, **DESCRIBE M2**
Before-after study (independent observations), **COMPARE2 A, B, H1, H2, H3**
Before-after study (paired observations), **PAIRSetc A, B, C, D1, D2, D3**
Before-after study (using changes in ratings), **COMPARE2 F2, PAIRSetc D1, D2**
Before-after study (using changes in ratings), sample size, **COMPARE2 S11**
Bias-adjusted kappa, **PAIRSetc A, B, C**
Binomial distribution, goodness of fit with, **DESCRIBE C**
Binomial goodness-of-fit test (any binomial probability), **PAIRS**
Binomial goodness-of-fit test (comparison with expected proportion) **DESCRIBE A**
Binomial success rate, **DESCRIBE C**
Birth-cohort effect (median polish procedure), **SMOOTH**
Bonferroni-like corrections (Finner, Haber and Hommel corrections), **ETCETERA C**
Bowley skewness coefficient, **DESCRIBE D**
Box-and-whisker diagrams, **DESCRIBE D**
Bradley-Blackwood test, **PAIRSetc D1, D2**

C
Calculator, **WHATIS calculator**
Capture-recapture method, **DESCRIBE I**
Case-control study, unmatched, **COMPARE2 C, G, LOGISTIC**
Case-control study, matched, **PAIRSetc A, C, D, E, H, J, L, LOGISTIC**
Centrifugality, **DESCRIBE D**
Change-point test, **DESCRIBE D**
Changes. based on before-after ratings, **COMPARE2 F2, PAIRSetc D1, D2**

Chi-square goodness-of-fit test, 2 x k table, **DESCRIBE C**
 Chi-square test, 2 x 2 table, **COMPARE2 A, B, C**
 Chi-square test, 2 x k table, **COMPARE2 F1**
 Chi-square tests for multiple comparisons in 2 x k table, **COMPARE2 F1**
 Chi-square, partitioning of, in 2 x k table, **COMPARE2 F1**
 Chi-square, partitioning of, in j x k table, **CHISQ**
 Chi-square, value corresponding to given P, or vice versa, **WHATIS P-value**
 Cluster-randomized trial, sample sizes, **COMPARE2 S1, S6**
 Cluster sample, required size, **DESCRIBE K**
 Cluster samples, comparison of binary data in, **COMPARE2 E, PAIRSetc A**
 Cluster samples, sample sizes for comparison, **COMPARE2 S1, S6**
 Clustering in space or time, **DESCRIBE C, H**
 Cochran Q test, **PAIRSetc F**
 Cochran-Armitage chi-square test for trend, **DESCRIBE B**
 Cochran-Orcutt procedure, **DESCRIBE B**
 Cohort study, unmatched, **COMPARE2 B, D, F, H, LOGISTIC, POISSON**
 Cohort study, matched, **PAIRSetc A, C, D, E, H, J, L, LOGISTIC**
 Combinations and permutations, **WHATIS calculator**
 Concordance coefficient, Kendall, **PAIRSetc I1**
 Concordance correlation coefficient, **PAIRSetc D1, D2, D3**
 Confidence intervals, **WHATIS C.I.**; also look up name of measure
 Confidence level for values above/below a given point, **WHATIS C.I.**
 Conover squared-rank test for difference between variances, **COMPARE2 H1**
 Correlation coefficient between values and ranks in sequence, **DESCRIBE D**
 Correlation coefficient, computation, **PAIRSetc D1, COMPARE2 F2**
 Correlation coefficient, test and confidence intervals for, **ETCETERA F1**
 Correlation coefficient, power and sample size for test, **ETCETERA F5**
 Correlation coefficient, based on Mantel-Haenszel procedure, **MANTELX**
 Correlation coefficients, comparison, **ETCETERA F2, F3**
 Correlation coefficients, partial and multiple, **ETCETERA F4**
 Correlation coefficients, pooling, **ETCETERA F2**
 Correlations, inter-item and item-total, in scale, **ETCETERA D**
 Covariance, analysis of, for comparing changes in two groups, **PAIRSetc D1, D2**
 Counts, heterogeneity of, **DESCRIBE D**
 Cox-Stuart test for trend, **DESCRIBE D**
 Cramer-von Mises test, **COMPARE2 F2, G, H1**
 Cressie-Read goodness-of-fit test, **DESCRIBE C**
 Cronbach's *alpha* (reliability-consistency) coefficient, **ETCETERA D**
 Cumulative frequency distribution, confidence interval for, **FIT**
 Cumulative odds ratio, **COMPARE2 F2, G**
 Cumulative rate, computed from age-specific rates, **DESCRIBE G**
 Cumulative survival proportions, **DESCRIBE F, COMPARE2 H3**
 Cumulative survival proportions, directly standardized, **COMPARE2 H3**
D
 D'Agostini-Pearson test for normality, **DESCRIBE D, COMPARE2 H2**
 Deciles, **DESCRIBE D**
 Departure from linear trend, **DESCRIBE B**
 Dersimonian-Laird estimates, **COMPARE2 A, B, C**
 Design effect, of cluster sample, **DESCRIBE J, COMPARE2 E**
 Diagnostic odds ratio, **DESCRIBE L3**
 Diagnostic tests, accuracy of, **DESCRIBE L1, L4, L5, PAIRSetc D1, D2, D3**
 Diagnostic tests, appraisal of, **DESCRIBE L1, L4**
 Diagnostic tests, combination of, **DESCRIBE L2**
 Diagnostic tests, comparison of, **DESCRIBE L2, L5**
 Diagnostic tests, meta-analysis, **DESCRIBE L3**
 Difference between paired values, **PAIRSetc D1, D3**
 Difference between proportions or risks, **COMPARE2 A, B, PAIRSetc A**
 Direct standardization, **DESCRIBE G**
 Distinguishability of categories, **PAIRSetc A, B, C**
 Donald-Donner procedure for comparing cluster samples, **COMPARE2 E**
 Durbin-Watson test, **DESCRIBE B**
E
 Edwards test for seasonal variation, **DESCRIBE E**

Effect sizes (standardized differences between means), combining, [COMPARE2 I](#)
 Epidemic curve, plotting of, using data, [DESCRIBE M1](#)
 Epidemic curve, plotting of, using models, [DESCRIBE M2](#)
 Epidemic models (Reed-Frost, SIR, SEIR), [DESCRIBE M2](#)
 Equivalence test for areas under ROC curve [DESCRIBE L5](#)
 Equivalence test for means, [COMPARE2 H2](#)
 Equivalence test for means, sample sizes for, [COMPARE2 S7](#), [PAIRSetc S7](#)
 Equivalence test for proportions, [COMPARE2 A](#), [PAIRSetc A](#)
 Equivalence test for proportions, sample sizes for, [COMPARE2 S2](#), [PAIRSetc S3](#)
 Exact tests, 2 x 2 table, power of, [COMPARE2 P1](#), [E](#)
 Exact tests, [COMPARE2 A, B, C, D, F1](#), [PAIRSetc A](#)
 Exponentially weighted moving averages, [SMOOTH](#)
 Expression solver (calculator), [WHATIS calculator](#)
F
 F (variance ratio) test, [COMPARE2 H2](#), [PAIRSetc I1](#)
 F, value corresponding to given P, or vice versa, [WHATIS P-value](#)
 Factorials, [WHATIS calculator](#)
 Fail-safe N, [COMPARE2 A, B, C, D, E, H3, I](#)
 False negative rate, [DESCRIBE L1](#)
 False positive rate, [DESCRIBE L1, L2, L3, L4](#)
 Finite population correction, [DESCRIBE A, D, K](#)
 Fisher exact tests, see Exact tests
 Fleiss-Everitt test for three ordered categories, [PAIRSetc C](#)
 Fligner-Policello robust rank test, [COMPARE2 H1](#)
 Forecasting efficiency, index of, [PAIRS](#)
 Freedman tests for seasonal variation, [DESCRIBE E](#)
 Frequency distribution, description of, [DESCRIBE D](#), [COMPARE2 H2](#)
 Friedman's two-way analysis of variance by ranks, [PAIRSetc I1](#)
 Funnel plot, skewed, [COMPARE2 A, B, C, D, I](#)
G
 Gain in certainty after diagnostic test, [DESCRIBE L1](#)
 Gamma (Goodman and Kruskal), [PAIRSetc C, D3](#)
 Gamma, in comparison of 2 samples, [COMPARE2 F2, G, H1](#), [PAIRSetc C, D3](#)
 Gehan-Wilcoxon test for comparison of survival, [COMPARE2 H3](#)
 Generalized odds ratio, [COMPARE2 F2, G](#), [PAIRSetc C](#)
 Geometric mean, [DESCRIBE D1](#)
 Geometric means, ratio of, [COMPARE2 H2](#)
 Gold standard, comparison with, [DESCRIBE L1](#), [PAIRSetc D1, D2, D3](#)
 Goodman and Kruskal's gamma, [COMPARE2 F2, G, H1](#), [PAIRSetc C, D3](#)
 Goodman and Kruskal's tau, 2 x k table, [COMPARE2 F1](#)
 Goodman and Kruskal's tau, j x k table, [CHISQ](#)
 Goodness of fit with expected distribution, [DESCRIBE C](#)
 Grubbs test for outliers, [DESCRIBE D](#)
 Guttman scale and coefficients of scalability and reproducibility, [ETCETERA D](#)
 Gwet's AC₁ statistic, [PAIRSetc A](#)
H
 Haber continuity correction, [COMPARE2 A, B, C](#)
 Haldane's large-table chi-square test, 2 x k table, [COMPARE2 F1](#)
 Haldane's large-table chi-square test, j x k table, [CHISQ](#)
 Harmonic-analysis procedure for appraisal of seasonal variation, [DESCRIBE E](#)
 Harmonic mean, [DESCRIBE D1](#)
 Hazard ratio, [COMPARE2 H3](#)
 Herd immunity, [DESCRIBE M2](#)
 Heterogeneity of correlation coefficients, [CORR](#)
 Heterogeneity of counts, [DESCRIBE D](#)
 Heterogeneity of season-specific slopes, [DESCRIBE D](#)
 Heterogeneity tests [COMPARE2 A, B, C, F2, G, H1, H2, I](#), [PAIRSetc A, C, D1 to D3, E, F, I1, J](#)
 Heterogeneity, measures of, [COMPARE2 A, B, C, F2, G, H1, H2, I](#), [PAIRSetc A, C](#)
 Hewitt's test for seasonal variation, [DESCRIBE E](#)
 Hollander's test for bivariate symmetry, [PAIRSetc D3](#)
 Hosmer-Lemeshow goodness-of-fit test, [LOGISTIC](#)
 Huber m-estimator, [DESCRIBE D](#)
I

Imputation of missing values in 2 x 2 table, **COMPARE2 A**
 Incidence rate, computed from survival data, **DESCRIBE F, COMPARE2 H3**
 Index of qualitative variation, **DESCRIBE C**
 Indirect standardization, **DESCRIBE H**
 Interobserver and intraobserver reliability, **PAIRSetc D1 to D3, I1, I2, M**
 Intraclass correlation coefficient in a cluster sample, **DESCRIBE J**
 Intraclass correlation coefficients, **COMPARE2 E, PAIRSetc D1, D2, I1, I2, M**
 Intraclass correlation coefficient, sample size, **PAIRSetc S6**
 Inverse sampling in comparative studies, **COMPARE2 A, B, C, PAIRSetc A**
 Inverse sampling in a descriptive study, **DESCRIBE A**
 Inverse sampling, sample size to find a given number of cases, **DESCRIBE K**

J

j x k table, **CHISQ, MANTELX**
 Jewell's low-bias estimate of odds ratio, **COMPARE2 A, B, C, PAIRSetc A, E, J**
 Jewell's low-bias estimate of ratio of proportions or risks, **COMPARE2 A, B**
 Jonckheere-Terpstra test for trend, **RANKCORR**

K

Kaplan-Meier life table procedure, **DESCRIBE F, COMPARE2 H3**
 Kappa, **PAIRSetc A, B, C, E, F, G, J, K**
 Kappa, in appraisal of screening or diagnostic test, **DESCRIBE L1**
 Kappa, sample size, **PAIRSetc S2**
 Kappa, weighted, **PAIRSetc C**
 Kendall concordance coefficient, **PAIRSetc I1**
 Kendall correlation coefficient (*tau*), **MANNWHIT, PAIRSetc C, D3**
 Kendall *tau* between ranks and values of numbers, **DESCRIBE D**
 Kendall *tau* in comparison of 2 samples, **COMPARE2 F2, G, H1, PAIRSetc C, D3**
 Kernel smoothing, **DESCRIBE B**
 Kernel smoothing, with log transformation, **SMOOTH**
 Knox test for space-time clustering, **DESCRIBE H**
 Kolmogorov-Smirnov test for a normal or lognormal distribution, **FIT**
 Kolmogorov-Smirnov test for an even distribution, **DESCRIBE D**
 Kolmogorov-Smirnov test for discrete data, **DESCRIBE C**
 Kolmogorov-Smirnov two-sample test, **COMPARE2 F2, G, H1**
 Kruskal-Wallis one-way analysis of variance by ranks (>2 samples), **MANNWHIT**
 Kullback-Leibler distances, **DESCRIBE L1, L4, L5**
 Kurtosis, **DESCRIBE D**

L

Lambda, **COMPARE2 A,B,C**
 Levene tests for equality of variances, **COMPARE2 H**
 Likelihood ratios for positive and negative tests, **DESCRIBE L1 to L5**
 Lilliefors test for normality, **DESCRIBE D, COMPARE2 H2**
 Linear regression, **DESCRIBE D, COMPARE2 F2, G**
 Linear regression, nonparametric, **DESCRIBE D**
 Log transformation, **DESCRIBE D, COMPARE2 H2, PAIRSetc D2, D3**
 Logistic regression analysis, conditional or unconditional, **LOGISTIC**
 Logit-rank test for effect of covariate on survival, **SURVIVAL**
 Logrank test for comparison of survival, **COMPARE2 H3**
 Low-cost optimal allocation, in stratified sample, **DESCRIBE K**

M

Mann-Kendall test for trend, **DESCRIBE D**
 Mann-Whitney test **COMPARE2 F2, G, H1**
 Mann-Whitney test for paired data, **PAIRSetc C**
 Mann-Whitney test, power, **COMPARE2 P3, PAIRSetc P2**
 Mann-Whitney test, sample sizes, **COMPARE2 S5, PAIRSetc S4**
 Mantel trend test, **DESCRIBE B**
 Mantel-Haenszel test, **COMPARE2 A, B, C, PAIRSetc E, J**
 Mantel-Haenszel test, extended, 2 x k table, **COMPARE2 F2, G, H1**
 Mantel-Haenszel test, extended, j x k table or numerical data, **MANTELX**
 Mantel-Haenszel test, power, **COMPARE2 P2**
 Mantel-Haenszel test, sample sizes, **COMPARE2 S3**
 McNemar test, **PAIRSetc A, C**
 McNemar test, extended, **PAIRSetc B**
 McNemar test, power, **PAIRSetc P1**

McNemar test, sample sizes, [PAIRSetc S1](#)
 Mean deviation from mean, [DESCRIBE D](#)
 Mean rank, [COMPARE2 H1](#)
 Mean square successive difference test, [DESCRIBE D](#)
 Mean, comparison with a hypothetical value, [DESCRIBE D](#)
 Mean, confidence interval, [WHATIS C.I.](#)
 Mean, robust estimators of, [DESCRIBE D](#)
 Means, difference between, [COMPARE2 H2](#), [PAIRSetc H, L](#)
 Means, difference between, required sample sizes, [COMPARE2 S6](#)
 Measures of impact of exposure, [COMPARE2 B, C, G](#), [PAIRSetc A](#)
 Median absolute deviation from median, [DESCRIBE D](#)
 Median polish procedure for two-way tables, [SMOOTH](#)
 Median rank, [COMPARE2 H1](#)
 Median survival times, ratio of, [COMPARE2 H3](#)
 Median, comparison with a hypothetical value, [DESCRIBE D](#)
 Median, [DESCRIBE D](#), [COMPARE2 H2](#)
 Medians, difference between, [COMPARE2 H1](#)
 Meta-analysis using available summary data, [COMPARE2 I](#)
 Meta-analysis, [COMPARE2 A, B, C, D, E, F1, F2, G, H1, H2, H3, I](#), [PAIRSetc A, B, C](#), [MANTELX](#)
 Meta-analysis, sensitivity analysis, [COMPARE2 I](#)
 Meta-analysis, studies of screening or diagnostic tests, [DESCRIBE L3](#)
 Mid-P exact tests, see [Exact tests](#)
 Minimization, [ETCETERA A6](#)
 Minimum significant change, [PAIRSetc I2](#)
 Misclassification, effect on appraisal of rate or proportion, [DESCRIBE A](#)
 Misclassification, effect on odds/risk ratio, [COMPARE2 M](#), [PAIRSetc M1, M2, M3](#)
 Missing values, imputation in 2 x 2 table, [COMPARE2 A](#)
 Moors kurtosis coefficient, [DESCRIBE D](#)
 Multiple comparisons, adjustment of P values, [ETCETERA C](#)
 Multiple comparisons in 2 x k table, [COMPARE2 F1](#)
 Multiple comparisons of categories, based on odds ratios, [PAIRSetc C](#)
 Multiple comparisons of correlation coefficients, [CORR](#)
 Multiple comparisons of means (Games-Howell procedure), [DIFFER](#)
 Multiple comparisons of rates or proportions, [DESCRIBE B](#), [PAIRSetc F](#)
 Multiple comparisons of samples, [PAIRSetc I](#)
 Multiple linear regression, sample sizes, [COMPARE2 S8](#)
 Multiple logistic regression, conditional or unconditional, [LOGISTIC](#)
 Multiple logistic regression, sample sizes, [COMPARE2 S4](#)
 Multiple Poisson regression, [POISSON](#)
 Multiple tests, adjustment of P-values for, [ADJUSTP](#)

N

Negative predictive value of a test, [DESCRIBE L1, L2](#)
 Neymann allocation, in stratified sample, [DESCRIBE K](#)
 Nominal-scale dependent variable, [COMPARE2 F](#), [PAIRSetc B](#)
 Nonparametric regression analysis, [DESCRIBE D](#)
 Nonparametric regression analysis, [PAIRSetc D3](#)
 Normality, tests for, [DESCRIBE D](#), [COMPARE2 H2](#)
 Number needed to treat, avoid/produce one case, [COMPARE2 B, C, D, H3](#), [PAIRSetc A](#)

O

Odds in favour of a higher value in one group, [COMPARE2 H1](#)
 Odds ratio, confidence interval, [WHATIS C.I.](#)
 Odds ratio, cumulative, [COMPARE F2, G](#)
 Odds ratio, diagnostic, [DESCRIBE L3](#)
 Odds ratio, for matched data, [PAIRSetc A, B, C, E, J](#)
 Odds ratio, generalized, [COMPARE2 F2, G](#), [PAIRSetc C](#)
 Odds ratio, in 2 x 2 table, [COMPARE2 A, B, C](#)
 Ordinal-scale dependent variable, [COMPARE2 F, G](#), [PAIRSetc C](#)
 Ordinal-scale variables, appraisal of association of, [PAIRSetc C, D3](#)
 Outliers, test for, [DESCRIBE D](#)
 Overall odds ratio, [COMPARE2 A, B, C, E](#)
 Overall's continuity correction, [CHISQ](#)

P

PABAK (prevalence-adjusted bias-adjusted kappa), [PAIRSetc A, B, C](#)

Page's test for trend, [PAIRSetc F, I](#)
 Paired *t* test, [PAIRSetc D1, D2, H, L](#)
 Paired *t* test, sample size, [PAIRSetc S5](#)
 Partitioning of chi-square in 2 x *k* table, [COMPARE2 F1](#)
 Percentage agreement, [PAIRSetc A, B, C](#) [DESCRIBE L1](#)
 Permutation (randomization) test, [COMPARE2 H1](#)
 Permutation test for matched pairs, [PAIRSetc C](#)
 Permutation test for paired replicates, [PAIRSetc D3](#)
 Permutations and combinations, [WHATIS calculator](#)
 Person-time, calculation of, [WHATIS imespan](#)
 Person-time rate, confidence interval for, [DESCRIBE A, WHATIS C.I.](#)
 Person-time rates, comparison of, [COMPARE2 D](#)
Phi, 2 x 2 or 2 x *k* table, [COMPARE2 A, B, C, F1](#)
 Pitman's test for equality of variances, [PAIRSetc D1, D2](#)
 Pococks harmonic analysis procedure, [DESCRIBE E](#)
 Poisson dispersion test, [DESCRIBE D](#)
 Poisson distribution, goodness of fit with, [DESCRIBE C](#)
 Poisson distribution, mean of, [DESCRIBE C](#)
 Poisson regression analysis, [POISSON](#)
 Poisson variate, confidence intervals for, [DESCRIBE H, WHATIS C.I.](#)
 Poisson variates, comparison of two numbers of events, [COMPARE2 D](#)
 Poisson variates, ratio of, confidence interval for, [WHATIS C.I.](#)
 Positive predictive value of a test, [DESCRIBE L1, L2](#)
 Post-test probabilities, [DESCRIBE L1, L2, L4, L5](#)
 Power of tests for comparison of independent samples, [COMPARE2 P1 to P6](#)
 Power of tests for comparison of matched samples, [PAIRSetc P1, P2, P3](#)
 Power of tests for regression and correlation coefficients, [REGPOWER](#)
 Predictive value of test result, [DESCRIBE L1, L2](#)
 Prevalence, confidence interval, [DESCRIBE A, I](#)
 Prevalence, estimation by capture-recapture method, [DESCRIBE I](#)
 Prevalence, estimation from cluster or stratified sample, [DESCRIBE J](#)
 Prevalence, estimation from screening test, [DESCRIBE L1](#)
 Prevalence, estimation from simple random sample, [DESCRIBE A](#)
 Prevalence-adjusted bias-adjusted *kappa*, [PAIRSetc A, B, C](#)
 Prevented or preventable fraction, [COMPARE2 B, C, G, PAIRSetc A](#)
 Proportion, comparison with hypothetical value, [DESCRIBE A, WHATIS C.I.](#)
 Proportion, confidence interval, [DESCRIBE A, WHATIS C.I.](#)
 Proportional allocation, in stratified sample, [DESCRIBE K](#)
 Proportions, difference between, [COMPARE2 A, B, PAIRSetc A, WHATIS C.I.](#)
 Proportions, sequence of, [DESCRIBE B](#)
P-value for given chi-square, *t*, *Z*, or *F*, or vice versa, [WHATIS P-value](#)
P-value, overall (pooled), [COMPARE2 I](#)
P-values, adjustment for multiple tests, [ETCETERA C](#)
Q
 Qstar (*Q**), [DESCRIBE L3](#)
 Quantiles (quartiles, quintiles, octiles, deciles), [DESCRIBE D](#)
R
r x *c* table, [CHISQ, MANTELX](#)
 Randles-Fligner-Policello-Wolfe test for symmetry, [FIT](#)
 Randles-Fligner-Policello-Wolfe test, [DESCRIBE D](#)
 Random decision (tossing a coin), [ETCETERA B7](#)
 Random effects model, [COMPARE2 A, B, C, I](#)
 Random numbers (tables), [ETCETERA B6](#)
 Random sampling, [ETCETERA B1, B2, B3, B4](#)
 Random sequence, [ETCETERA A7, B5](#)
 Randomization test, [COMPARE2 H1](#)
 Randomization, [ETCETERA A1, A2, A3, A4, A5](#)
 Randomness, tests for, [DESCRIBE D](#)
 Rank correlation between ordinal-scale variables, [PAIRSetc C, D3](#)
 Rank correlation between ranks and values of numbers, [DESCRIBE D](#)
 Rank correlation coefficients, partial, [RANKCORR](#)
 Rank correlation in comparison of two samples, [PAIRSetc C, D3](#)
 Rao-Scott procedure for comparing cluster samples, [COMPARE2 E](#)

Ratchet circular scan test for seasonal variation, **DESCRIBE E**
 Rate difference, **COMPARE2 D, WHATIS C.I.**
 Rate of homogeneity. in cluster sample, **DESCRIBE J**
 Rate ratio, **COMPARE2 D, WHATIS C.I.**
 Rate, comparison with hypothetical value, **DESCRIBE A**
 Rate, confidence interval, **DESCRIBE A, WHATIS C.I.**
 Rates, sequence of, **DESCRIBE B**
 Rates, person-time, comparison of, **COMPARE2 D**
 Ratio of paired values, **PAIRSetc D2, D3**
 Ratio of proportions or risks, **COMPARE2 A, B**
 Ratios of counts. sequence of, **DESCRIBE B**
 Recurrence, chances of, **DESCRIBE A**
 Reed-Frost epidemic model, **DESCRIBE M2**
 Regression analysis, **DESCRIBE D, COMPARE2 F2, G**
 Regression analysis, Mantel-Haenszel procedure, **MANTELX**
 Regression analysis, nonparametric, **PAIRSetc D3**
 Regression analysis, sequence of rates or proportions, **DESCRIBE B**
 Regression asymmetry test for skewed funnel plot, **COMPARE2 A, B, C, D, I**
 Regression equations, reciprocal/log-root/exponential/power/quadratic, **PAIRS**
 Regression to the mean, estimated effect of, **PAIRSetc D5**
 Relative difference between proportions, **PAIRSetc A**
 Relative survival curve, **LIFETAB**
 Reliability of numerical observations, **PAIRSetc D1 to D3, I1, I2, M**
 Reliability of ratings, using kappa, **PAIRSetc A, B, C, E, F, G, J, K**
 Reliability of scale (*alpha* coefficient), **ETCETERA D**
 Repeatability, coefficient of, **PAIRSetc D1, D2, H, I1, I2, M**
 Reproducibility, Guttman coefficient of, **ETCETERA D**
 Replicate measurements, **PAIRSetc D1, D2, D3, I1, I2, M**
 Residuals, adjusted (chi-square test), **CHISQ**
 Redit analysis, **COMPARE2 F**
 Risk ratio, **COMPARE2 A, B, PAIRSetc A, WHATIS C.I.**
 Risk, comparison with hypothetical value, **DESCRIBE A, WHATIS C.I.**
 Risk, computed from age-adjusted (cumulative) rate, **DESCRIBE G**
 Risk, confidence interval, **DESCRIBE A, WHATIS C.I.**
 Risk difference, **COMPARE2 A, B, PAIRSetc A**
 Risk marker, appraisal of, **DESCRIBE L1**
 Robust estimators of mean, **DESCRIBE D**
 ROC curve and area under ROC curve, **DESCRIBE L4, L5**
 ROC curve, in meta-analysis of screening or diagnostic tests, **DESCRIBE L3**
 Rosner's test, **PAIRSetc H, L**
 Running medians, **DESCRIBE D**
 Runs test for serial correlation, **DESCRIBE B**
 Runs tests for randomness, **DESCRIBE D**
S
 Sample, random, selection of, **ETCETERA B1, B2, B3, B4**
 Sample size for descriptive study, **DESCRIBE K**
 Sample size for estimating proportion, rate, or mean, **DESCRIBE K**
 Sample size for regression and correlation analysis, **REGPOWER**
 Sample size to find a given number of cases, **DESCRIBE K**
 Sample sizes for comparison of cluster samples, **COMPARE2 S1, S6**
 Sample sizes for comparison of matched samples, **PAIRSetc S1 to S7**
 Sample sizes for comparison of two independent samples, **COMPARE2 S1 to S11**
 Scalability, coefficient of, **ETCETERA D**
 Scale, internal consistency and discriminatory power of, **ETCETERA D**
 Screening and diagnostic test, optimal cut-point for, **DESCRIBE L4, L5**
 Screening tests, accuracy of, **DESCRIBE L1, L4, L5, PAIRSetc D1, D2, D3**
 Screening tests, appraisal of, **DESCRIBE L1, L4, L5**
 Screening tests, combination of, **DESCRIBE L2**
 Screening tests, comparison of, **DESCRIBE L2, L5**
 Screening tests, meta-analysis, **DESCRIBE L3**
 Seasonal Mann-Kendall test, **DESCRIBE D**
 Seasonal variation, appraisal of, **DESCRIBE E**
 Seasonal variation, control of, **DESCRIBE D**

SIR epidemic model, **DESCRIBE M2**
 Sen estimator of slope, **DESCRIBE D**
 Sensitivity analysis in meta-analysis, **COMPARE2 I**
 Sensitivity of a measure or test, **DESCRIBE L1 to L5**
 Sensitivity, chance-corrected, **DESCRIBE L1, L2**
 Sensitivity test in meta-analysis, **COMPARE2 I**
 Sequence of numbers, appraisal of, **DESCRIBE D**
 Sequence of rates or proportions or ratios, appraisal of, **DESCRIBE B**
 Serial correlation of residuals, **DESCRIBE B**
 Shannon's index of the diversity of distribution, **COMPARE2 F1**
 Shapiro-Francia W' test for normality, **FIT**
 Shoemaker's test for comparing two variances, **COMPARE2 H2**
 Sign test, **PAIRS**
 SIR epidemic model, **DESCRIBE M2**
 Skewed funnel plot, **COMPARE2 A, B, C**
 Skewness, **DESCRIBE D**
 Smoothing of curve, **DESCRIBE D**
 Smoothing of curve, exponentially weighted moving averages, **SMOOTH**
 Smoothing of curve, kernel smoothing, **DESCRIBE B**
 Smoothing of curve, kernel smoothing, with log transformation, **SMOOTH**
 Smoothing of curve, median and quartile paths, **SMOOTH**
 SMR (standardized morbidity or mortality ratio), **DESCRIBE H**
 Somers' D, **PAIRSetc C, D3**
 Space-time clustering, Knox test, **DESCRIBE H**
 Spearman correlation coefficient (ρ) **PAIRSetc C, D3**
 Spearman ρ between ranks and values of numbers, **DESCRIBE D**
 Specificity of a measure or test, **DESCRIBE L1 to L5**
 Specificity, chance-corrected, **DESCRIBE L1, L2**
 St Laurent's gold-standard correlation coefficient, **PAIRSetc D1, D2**
 Standard deviation, confidence interval for, **WHATIS C.I.**
 Standard error of measurement, **PAIRSetc D1, D2, I2**
 Standard populations, world, European and African, **DESCRIBE G**
 Standardization, direct, **DESCRIBE G**
 Standardization, indirect, **DESCRIBE H**
 Standardized morbidity or mortality ratio, **DESCRIBE H**
 Standardized odds ratios, **COMPARE2 C**
 Stratified sample, for estimation of prevalence, **DESCRIBE J**
 Stratified sample, required size, **DESCRIBE K**
 Strickland-Lu test for difference between changes, **COMPARE2 F2**
 Strickland-Lu test, sample sizes for, **COMPARE2 S11**
 Stuart-Maxwell test for marginal heterogeneity, **PAIRSetc B, C**
 Summary ROC curve, **DESCRIBE L3**
 Survival curve, **DESCRIBE F**
 Survival curve, relative to expectation, **LIFETAB**
 Survival periods, calculation of, **WHATIS timespan**
 Survival time, **DESCRIBE F**
 Survival time, effects of covariates on, **SURVIVAL**
 Survival time, mean and median, **DESCRIBE F, COMPARE2 H3**
 Survival times, comparison of, **COMPARE2 H3**
 Survival times, comparison of, sample sizes for, **COMPARE2 S10**
 Symmetry test, **PAIRSetc B**
 Synergism, appraisal of, **ETCETERA E**
 Synergism, appraisal of (median polish procedure), **SMOOTH**
T
 t, value corresponding to given P, or vice versa, **WHATIS P-value**
 Theil's uncertainty coefficient U, **COMPARE2 F1**
 Time intervals, **WHATIS timespan**
 Time-to-event data, **DESCRIBE F, COMPARE2 H3**
 Trend of rate ratios, **COMPARE2 D**
 Trend of rates or proportions, **DESCRIBE B**
 Trend of three or more samples, Cuzick test, **MANNWHIT**
 Trend of three or more samples, Pee-Freedman test, **MANNWHIT**
 Trend, **DESCRIBE D**

Trend, Jonckheere-Terpstra test, *RANKCORR*

Trial, unmatched controls, *COMPARE2 B, D, F, H, LOGISTIC, POISSON*

Trial, matched controls, *PAIRSetc A, C, D, E, H, J, L, LOGISTIC*

Trimmed mean, *DESCRIBE D*

t-test, *COMPARE2 H2*

t-test, power, *COMPARE2 P4*

U

Up-and-down runs test for randomness, *DESCRIBE D*

Upton continuity correction, *COMPARE2 A,B,C*

V

Validity study, using "gold standard", *DESCRIBE L1, PAIRSetc D1 to D3*

Van Elteren procedure, *COMPARE2 F2, G, H1*

Variance inflation factor, in cluster sample, *DESCRIBE J, COMPARE2 E*

Variance ratio, confidence interval for, *COMPARE2 H2*

Variance, confidence interval for, *WHATIS C.I.*

Variances, comparison of, *COMPARE2 H2, PAIRSetc I1*

W

Walter test for numerical data, *PAIRSetc H ,L*

Walter test for binary data, *PAIRSetc E, J*

Wilcoxon rank-sum test, *COMPARE2 F2, G ,H1*

Wilcoxon signed-rank test of symmetry, *DESCRIBE D*

Wilcoxon signed-rank test, *PAIRSetc C , D3*

Y

Yates continuity correction, *COMPARE2 A, B, C*

Youden's index, *DESCRIBE L1, L2, L4, L5*

Yule's Q , *COMPARE2 A, B ,C*

Z

Z, value corresponding to given P, or vice versa, *WHATIS P-value*
